

International Society for Sexual Medicine's Guidelines for the Diagnosis and Treatment of Premature Ejaculation

Stanley E. Althof, Ph.D., Carmita H.N. Abdo, M.D., Ph.D., John Dean, M.D., Geoff Hackett, M.D., Marita McCabe, Ph.D., Chris G. McMahon, M.D., Raymond C. Rosen, Ph.D., Richard Sadovsky, M.D., Marcel D. Waldinger, M.D., Ph.D., Edgardo Becher, M.D., Gregory A. Broderick, M.D., Jacques Buvat, M.D., Irwin Goldstein, M.D., Amr I. El-Meliegy, M.D., Francois Giuliano, M.D., Ph.D. Wayne J.G. Hellstrom, M.D., Luca Incrocci, M.D., Emmanuelle Jannini, M.D., Kwangsung Park, M.D., Sharon Parish, M.D., Hartmut Porst, M.D., David Rowland, Ph.D., Robert Seagraves, M.D., Ph.D., Ira Sharlip, M.D., Chiara Simonelli, Ph.D., Hui Meng Tan, M.D.

Introduction

Over the past 20 years our knowledge of premature ejaculation (PE) has significantly advanced¹. Specifically, we have witnessed substantial progress in understanding the physiology of ejaculation²⁻⁸, clarifying the real prevalence of PE in population-based studies⁹⁻¹³, reconceptualizing the definition and diagnostic criterion of the disorder¹⁴, assessing the psychosocial impact on patients and partners¹⁵, designing validated diagnostic and outcome measures¹⁶, proposing new pharmacologic strategies and examining the efficacy, safety and satisfaction of these new and established therapies¹⁷⁻²¹. Given the abundance of high level research it seemed like an opportune time for the International Society for Sexual Medicine (ISSM) to promulgate an evidenced-based, comprehensive and practical set of clinical guidelines for the diagnosis and treatment of premature ejaculation.

Treatment guidelines provide specific recommendations about clinical interventions. They tend to be condition or treatment specific and typically disorder based²². Guidelines are useful to aid clinicians in making decisions about the management of individual patients. Attributes for good primary care guidelines include: 1) beginning with a systemic review of the relevant literature, 2) taking into consideration the level of methodological rigor and clinical sophistication of the research supporting the intervention, 3) proposing strong recommendations only when convincing patient-oriented evidence exists, 4) making recommendations that are based on patient-oriented outcomes rather than disease oriented outcomes, 5) considering the ability of health care professionals to use, and of patients to accept, the recommendations offered, 6) having a transparent development process, 7) identifying potential conflicts of interest, 8) ongoing prospective validation (adoption of guideline has been shown to improve patient-oriented outcomes in a real world setting), and 9) flexibility in various clinical situations^{22, 23}.

All evidence is not created equal and there are several excellent systems to evaluate evidence and propose recommendations. Guidelines should offer health care professionals (HCPs) reasonable levels of confidence about making decisions to support clinical activities. The two

most widely accepted systems for evaluating treatment recommendations are the Oxford Centre of Evidence-Based Medicine and the Strength of Recommendation Taxonomy (SORT) developed by the Association of Family Practitioner^{24, 25}. Prior to disseminating recommendations each system carefully scrutinizes study design, quantity, quality and consistency of the evidence as well as the magnitude of the effect, bias or limitations of the evidence and patient values.

We were able to identify three sets of clinical practice guidelines for the diagnosis and treatment of premature ejaculation. They are the American Urological Association's 2004 Guidelines for the Pharmacologic Treatment of Premature Ejaculation, the European Association of Urology Guidelines on Male Sexual Dysfunction: Erectile Dysfunction and Premature Ejaculation, Updated 2009 and the Practice Guidelines for the Pan Arab Society of Sexual Medicine (Disorders of Ejaculation)²⁶⁻²⁹. It was our opinion that the existing guidelines were not sufficiently comprehensive, failed to adequately address both psychological as well as medical interventions and that essential new evidence that has come to light is not included in these documents. Therefore we undertook to develop a contemporary set of practical guidelines primarily targeting front line clinicians, and secondarily sexual medicine specialists.

Guideline Development Process

The ISSM determined that the utility of the guideline would be enhanced if it were developed by an independent panel of experts, from different professions and clinical disciplines, and from different cultural and national backgrounds. The guideline and related resources should be practical and clinically relevant, as well as authoritative, as busy health care professionals (HCPs) and patients need guidance to deal with clinical problems posed by PE on a day-to-day basis. The guideline needed to be accessible for clinicians from all disciplines, including family physicians and other specialists without expertise in sexual medicine.

In September, 2009 the ISSM PE Guidelines Committee met in London for three days. The 26 committee members were selected by peer recommendation and further vetted to provide a diversity of discipline, balance of opinion, knowledge, gender and geography. These 22 men and 4 women included most of the world's highly recognized experts on PE and comprised ten urologists, five psychologists, three psychiatrists, two endocrinologists, two primary care physicians, one sexual medicine physician, one genitourinary physician, one internal medicine physician, and one radiation oncologist. The committee was chaired by Stanley Althof, Ph.D. and the meeting facilitated by ISSM President, John Dean, M.D.. All members were required to declare in advance any potential conflicts of interest with their participation in the committee's work.

A comprehensive review of scientific literature on PE was conducted by ISSM research assistants under the supervision and guidance of the Chairman. Selected original articles were circulated to committee members prior to meeting. Committee members were invited to review all available evidence, not just the articles provided by the ISSM researchers, and prepare a presentation for the meeting on specific topics related to the guideline. The committee employed evidence-based medicine grading as the means of integrating individual clinical expertise with the best available external clinical evidence from systematic research.

Where available, evidence of important outcomes was collated for each sub-population. Quality of evidence, and the strength of any recommendation was graded using the Oxford Centre of Evidence-Based Medicine system³⁰. The relevant evidence-based recommendations of the Committee are summarized in Table 1 and have been published in the Journal of Sexual Medicine³¹.

The meeting was supported by an unrestricted grant from Johnson and Johnson, the manufacturer of dapoxetine. However, ISSM required complete independence from industry during the development of the guideline and related resources. There were no industry representatives at the meeting and there was no attempt by industry to influence any part of the development or writing process at any time.

Definitions of Premature Ejaculation

Several definitions for premature ejaculation (PE) exist having been crafted by various professional organizations and/or individuals^{11, 29, 32-36}(see Table 2). Most of these definitions include the subtypes of lifelong and acquired (PE symptoms beginning after a period of normal ejaculatory function). The major criticisms of the extant definitions included their failure to be evidenced-based, lack of specific operational criteria, excessive vagueness, and reliance on the subjective judgment of the diagnostician. Nonetheless, three common constructs underlie most definitions of PE: 1) a short ejaculatory latency; 2) a lack of perceived self-efficacy or control about the timing of ejaculation; and 3) distress and interpersonal difficulty (related to the ejaculatory dysfunction).

Because of the discontent with the existing definitions of PE, as well as pressure from the regulatory agencies concerning the inadequacy of the PE definitions, the ISSM convened in 2007 a meeting of experts to review the evidence-based literature and to develop a definition grounded in clearly definable scientific criteria¹⁴. After carefully reviewing the literature, the Committee proposed that **lifelong PE** is,

“a male sexual dysfunction characterized by ejaculation which always or nearly always occurs prior to or within about one minute of vaginal penetration, and the inability to delay ejaculation on all or nearly all vaginal penetrations, and negative personal consequences, such as distress, bother, frustration and/or the avoidance of sexual intimacy.” (LOE 1a)

The definition applies only to intravaginal sexual activity. It does not define premature ejaculation in the context of other sexual behaviors or men having sex with men. The Committee concluded that there is insufficient information available to extend the definition of PE to these other groups. **Additionally, the Committee concluded that there are insufficient published objective data to propose a new evidence-based definition of acquired PE, although it believed the proposed criterion for lifelong PE might be applied to acquired PE as well. (LOE 5d)**

Anteportal ejaculation is the term for men who ejaculate prior to vaginal penetration and is considered the most severe form of premature ejaculation. Such men/couples typically present when they are having difficulty conceiving children. It is estimated that between 5% and 20 % of lifelong PE-men suffer from anteportal PE³⁷.

Although not evidenced based, Waldinger proposed two additional “subtypes” for men who are distressed about their ejaculatory function but do not meet the ISSM criterion for PE³⁶. These subtypes should be considered provisional; however, we thought it was important to include them because they accurately characterize many men who do not qualify for the diagnosis of PE and are asking for help. We believe these categories may help health care professionals (HCPs) address these men’s concerns.

These two subtypes are termed Natural Variable PE and Premature Like Ejaculatory Dysfunction. Natural variable PE is characterized by early ejaculations which occur irregularly and inconsistently with some subjective sense of diminished control of ejaculation. This subtype is not considered a sexual dysfunction or psychopathology but rather a normal variation in sexual performance. Premature-like ejaculatory dysfunction is characterized by: 1) subjective perception of consistent or inconsistent rapid ejaculation during intercourse; 2) preoccupation with an imagined early ejaculation or lack of control of ejaculation; 3) actual intravaginal ejaculation latency time in the normal range or even of longer duration (i.e. an ejaculation that occurs after 5 minutes); 4) ability to control ejaculation (i.e. to withhold ejaculation at the moment of imminent ejaculation) may be diminished or lacking and; 5) the preoccupation is not better accounted for by another mental disorder³⁸. **(LOE 5d)**

Prevalence

Reliable information on the prevalence of lifelong and acquired PE in the general male population is lacking. Confounding accurate prevalence estimates are the competing and varying definitions of PE and the manner in which prevalence data was gathered (population-based, self-report, or clinician based). Local and regional variations should be considered in the context of different cultural, religious, and political influences. Additionally, prevalence may vary across different demographics, including geography, ethnicity and social status¹¹.

Based on patient self report, PE is routinely characterized as the most common male sexual dysfunction. Prevalence data derived from patient self report will be appreciably higher than prevalence estimates based on clinician diagnosis utilizing the more conservative ISSM definition of PE. The following studies demonstrate the varying prevalence estimates ranging from 30% down to 3%. Data from The Global Study of Sexual Attitudes and Behaviors (GSSAB), an international survey investigating the attitudes, behaviors, beliefs, and sexual satisfaction of 27,500 men and women aged 40-80 years, reported the global prevalence of PE (based on subject self-report) to be approximately 30% across all age groups^{9, 12}. Perception of “normal” ejaculatory latency varied by country and differed when assessed either by the patient or their partner³⁹. A core limitation of the GSSAB survey stems from the fact that the youngest participants were aged 40 years, an age when the incidence of premature ejaculation might be different from younger males.⁴⁰

Contrary to the GSSAB study, the Premature Ejaculation Prevalence and Attitude Survey found the prevalence of PE among men aged 18 to 70 to be 22.7%¹⁰. PE was categorized by subject self-report. Prevalence was similar across countries (24.0% in the United States,

20.% in Germany and Italy) and age groups. Similar results by age groups were also found by Brazilian Sexual Life Study (BSLS), and PE was reported in 25.8% of 3332 Brazilian men⁴¹.

A Canadian study, using a study-specific definition of PE, based largely on the DSM III criteria (those who had fair/to poor control over their ejaculation and claim that time to climax was a problem for them or their partner), classified 16% of males aged 18 to 60 years and above as experiencing PE⁴². Consistently lower rates of PE were reported by female partners, ranging between 9% and 14%. No Canadian regional differences were noted. When analyzed by age group, there was only minor variation, with the highest prevalence in the male respondents seen in the age 55-64 years age group (18%), while the lowest rate was observed in the 18-24 years age group (12%).

Applying only the time parameter (IELT approximately 1 minute) of the ISSM definition of PE to a population based cohort of 500 men with stopwatch measured IELT's, only 1%-3%, would be eligible for the diagnosis^{40, 43}. These men were not given measures to assess control or distress. **The 1% - 3% prevalence is considerably different from the previous prevalence estimates of 20% - 30% based on self-report or DSM-III diagnosis. The lower prevalence estimates are however more consistent with the numbers of men who present for treatment of PE. (LOE 3b)**

Prevalence of Premature Ejaculation in Clinical Practice

Premature ejaculation in clinical practice is frequently a self-reported complaint, making it difficult to appreciate its real epidemiology. In addition, in some men/couples premature ejaculation is diagnosed on the basis of distress rather than as an objective symptom. Another problem is the relative inconstancy of the symptom in many patients. The real prevalence is difficult to assess in clinical practice⁴⁰.

Very few PE sufferers seek help (9%); of those who seek help, 91.5% report little or no improvement as a result of seeking treatment; and in most cases (81.9%), it is the sufferer himself who seeks to initiate the conversation¹⁰. Reporting on practice patterns of urologists, Shindel et al notes that respondents saw one new PE patient per week and 26% prescribed on-demand SSRI's, 22% daily SSRIs and 11% topical anesthetics⁴⁴.

Average Ejaculatory Latency

Normative multinational (Netherlands, United Kingdom, United States, Spain, and Turkey) reports of IELT been only recently been published¹¹. **The median IELT was 5.4 minutes (range, 0.55–44.1 minutes) and the distribution of the IELT in all five countries was positively skewed.** The median IELT decreased significantly with age, from 6.5 minutes in the 18–30 years group, to 4.3 minutes in the group older than 51 years. Median IELT varied between countries, with Turkey having the lowest IELT. The median IELT value was independent of condom use or circumcision status (except in Turkey). A similar study conducted a few years later reported congruent results with a median IELT of 6 (range 0.1-15.2 minutes)⁴³. **(LOE 2a)**

Etiology

The distinction between the four PE subtypes or syndromes (lifelong, acquired, Natural Variable and Premature Like Ejaculatory) underscores that there is no one particular pathophysiology of PE. Attempts to explain the etiological factors for PE have included a diverse range of biological and psychological factors.

In 1943, Bernhard Schapiro first distinguished between lifelong and acquired⁴⁵. However, Schapiro's descriptive classification failed to distinguish between specific etiological factors that were associated with both forms of the dysfunction. Many in the field assumed that both forms were primarily psychologically or interpersonally based³²

Over the past two decades several studies have suggested that lifelong and acquired PE may be caused by somatic disorders and/or neurobiological disturbances. Previously it was thought that the condition was primarily psychologically or interpersonally based. Examples of these biological factors include: hypersensitivity of the glans penis⁴⁶, a higher cortical representation of the pudendal nerve⁴⁷, disturbances in central serotonergic neurotransmission^{48, 49}, erectile difficulties⁵⁰, prostatitis^{51, 52}, detoxification from prescribed drugs (e.g. raxoxetine⁵³) or recreational drugs^{54, 55}, chronic pelvic pain syndrome⁵⁶ and thyroid disorders^{57, 58}. It is noteworthy that **none of these "organic" etiologies has yet been supported by robust and large scale studies.**

Recent studies have suggested that in some men neurobiological and genetic variations could contribute to the pathophysiology of lifelong PE, as defined by the ISSM criteria and that the condition may be maintained and heightened by psychological/environmental factors^{59, 60}. In animal experiments serotonin receptors have been implicated in the etiology of lifelong PE and partly in acquired PE. Waldinger et al hypothesized that lifelong early ejaculation in humans may be explained by either a hyposensitivity of the 5-HT_{2C} and /or hypersensitivity of the 5-HT_{1A} receptors⁶¹. They hypothesized that men with a low 5-HT neurotransmission and probable 5-HT_{2C} receptor hyposensitivity may have their ejaculatory threshold genetically set at a lower point and ejaculate quickly with minimal stimulation. Additionally, in rapidly ejaculating rats, it has been recently demonstrated that the spinal command of ejaculation differs when compared to "normal" rats⁶². **Serotonin dysregulation as an etiological hypothesis for PE explains only a small percentage (2-5%) of complaints of PE in the general population⁵⁹.**

Genetics of PE

In his classic paper, Bernhard Schapiro noted that some family members of men with PE also have PE⁴⁵. Many years later Waldinger et al hypothesized that both the IELT and lifelong PE for some men are genetically determined⁴⁸. Supporting this hypothesis was Waldinger et al's report of a high prevalence of lifelong PE (defined in terms of and IELT of less than 1 minute) among first degree male relatives of Dutch men with lifelong PE⁶³.

Models to explain premature and delayed ejaculation were developed by Jern based on examination of 1196 Finish male twins between the ages of 33 to 43⁶⁰. The premature ejaculation model suggested a moderate additive genetic variance of 28%, with no shared environmental variance (0%) and 72% non-shared environmental variance. One possible

interpretation of these findings is that genetic influences may create a diathesis or predisposition in some men to ejaculate prematurely.

The first DNA study on PE was performed by Janssen et al in 89 Dutch men with lifelong PE, in whom IELT was measured by a stopwatch⁶⁴. The data were compared to a cohort of the mentally and physically healthy Dutch Caucasian men. He demonstrated an association of the 5-HTLPR gene polymorphism and the duration of the IELT. Men with lifelong PE, defined as IELTs of less than 1 minute, and with the LL genotype ejaculated in a 100% shorter time than PE men with the SL and SS genotype. Additionally, the study showed that there was no difference in the prevalence of the LL, SL and SS genotypes in men with lifelong PE compared to their prevalence in the general male Dutch population. Given that the distribution of the polymorphism is similar in lifelong premature ejaculators and mentally and physically healthy Dutch men **this study also lends support to a predisposition/diathesis model rather than the conclusion that genetic influences underlie all men with lifelong premature ejaculation.**

In contrast to the study of Janssen et al,⁶⁴ a study in Turkish men⁶⁵ and Iranian men⁶⁶ showed a higher prevalence of SS genotypes in men with lifelong PE. However, the latter studies have been criticized on their methodology and/or use of statistics^{67, 68}. **(LOE 2a)**

Thyroid Hormones

Data from animal studies suggest anatomic and physiologic interactions between brain dopamine and serotonin systems and the hypothalamic-pituitary thyroid axis^{69, 70}. **The data linking thyroid hormones and ejaculatory dysfunction is inconsistent.** Corona et al. reported a significant correlation between PE and suppressed TSH values in a selected population of andrological and sexological patients.⁷¹ Carini et al reported that after normalizing thyroid function in hyperthyroid men the prevalence of PE fell from 50% to 15%^{57, 58, 72}. However, contradicting this data is the failure to find an association between thyroid hormones and PE in a large cohort of men with lifelong PE⁷³. **(LOE 2a)**

Prostatitis

Acute and chronic lower urogenital infection, prostatodynia, or chronic pelvic pain syndrome (CPPS) is associated with ED, PE and painful ejaculation. The relationship between chronic prostatitis, CPPS and premature ejaculation is supported by several recently published studies which focus more on epidemiology and largely ignore treatment. Most of these have are limited by poor study design including inconsistent or absent methodologies of microbiological diagnosis of prostatitis and the lack of a validated questionnaire for combined evaluation of chronic prostatitis and sexual dysfunction.

Twenty-six to seventy-seven percent of men with chronic prostatitis or chronic pelvic pain syndrome (CPPS) report experiencing PE^{56, 74, 75}. Prostatic inflammation and chronic bacterial prostatitis have been reported as common findings in men with acquired PE^{51, 52, 76}. Considering the role of the prostate in the ejaculatory mechanism, a direct influence of the local inflammation in the pathogenesis of some cases of acquired PE seems likely⁷⁷. **The exact pathophysiology of the link between chronic prostatitis, ED and PE is unknown. (LOE 3a)**

Psychological Factors

There are a range of psychological factors that may precipitate or maintain PE. These factors can be divided into predisposing or historical factors (e.g., sexual abuse, attitude toward sex in the home), individual psychological factors (e.g., body image, depression, performance anxiety, alexithymia), or relationship factors (e.g., intimacy, anger)⁷⁸⁻⁸⁰. There has been limited research that investigated the direction of the relationship between the above variables and PE. Most studies are cross-sectional in nature, and associations between the variables have been identified, rather than causative relationships. Care must be exercised in interpreting these relationships, as it is possible that either the psychological factors led to the PE, or vice versa. From a clinical perspective, it is likely that there is a reciprocal relationship occurring, where one condition has led to problems in other areas, and these have, in turn, exacerbated the original condition. For example, performance anxiety may lead to PE, which then further exacerbates the original performance anxiety. **(LOE 5d)**

Quality of Life Impact of PE on the Man and Partner

Rosen and Althof reviewed 11 observational, non-interventional studies from 1997 through 2007 that report on the psychosocial and quality of life consequences of PE on the man, his partner and the relationship¹⁵. These studies employed different methodologies, outcome measures and consisted of both qualitative and quantitative investigations. All studies consistently confirmed a high level of personal distress reported by men with PE and their female partners. The negative impact on single men with PE may be greater than on men in relationships as it serves as a barrier to seeking out and becoming involved in new relationships⁸¹. Men with PE have significantly lower scores on self-esteem and self-confidence than non-PE men and one-third of men with PE report anxiety connected to sexual situations⁸². It may seem obvious but PE has a direct negative influence on women's sexual experience^{10, 83}. **Both men and their partners affirm negative effects and interpersonal difficulty related to their PE and an overall reduction in their quality of life^{15, 82, 84}. (LOE 1a-3a)**

Should Clinicians Specifically Screen for PE

Screening involves testing of a symptomless population in order to detect cases of a disease at an early stage. Clearly, men affected by PE are not symptomless and, for PE, case-finding might be a better term.

The committee agreed that there was inadequate evidence to recommend screening or case-finding for PE, either in a general population or in most sub-population. However, it is recommended that men with ED be screened for PE (LOE 5d).

Improvements in public awareness of sexual health issues, including PE, are required, so that individuals affected by sexual concerns are aware of options and interventions open to them. Physicians have an important role to play in sexual health education and PE should be included within sexual health education programs.

Assessment of PE

History

Patients expect clinicians to inquire about their sexual health⁸⁵. Often patients are too embarrassed, shy, and uncertain if sexual complaints belong in the HCP's office⁸⁶. Inquiry into sexual health gives patients' permission to discuss their sexual concerns and also screens for associated health risks (e.g. cardiovascular risk and ED).

Table 3 lists recommended and optional questions that patients who complain of PE should be asked⁸⁷. The recommended questions establish the diagnosis and direct treatment considerations and the optional questions gather detail for implementing treatment. Finally, the committee recommends that the HCPs take a medical and psychosocial history (LOE 5d)

Figure 1 is a flowchart devised by Rowland et al, detailing the assessment and treatment options for subjects complaining of premature ejaculation¹.

Assessment of Ejaculatory Latency

Stopwatch Assessment of Ejaculatory Latency (IELT)

Stopwatch measures of IELT are widely used in clinical trials and observational studies of PE, but have not been recommended for use in routine clinical management of PE. Despite the potential advantage of objective measurement, stopwatch measures have the disadvantage of being intrusive and potentially disruptive of sexual pleasure or spontaneity. More recently, studies have indicated that patient or partner self-report of ejaculatory latency correlate relatively well with objective stop-watch latency and might be useful as a proxy measure of IELT⁸⁸⁻⁹¹. **Since patient self-report is the determining factor in treatment seeking and satisfaction, it is recommended that self-estimation by the patient and partner of ejaculatory latency be accepted as the method for determining IELT in clinical practice. (LOE 2b)**

Physical Examination

For lifelong PE, a physical examination is highly advisable but not mandatory and should be conducted in most if not all patients. Some patients find it reassuring for the physician to examine them.

For acquired PE a targeted physical examination is mandatory to assess for associated/causal diseases such as ED, thyroid dysfunction and prostatitis. (LOE 5d)

Use of Assessment Instruments or Required Stopwatch Assessments

Standardized assessment measures for premature ejaculation include the use of validated questionnaires and patient reported outcome (PRO) measures, in addition to stopwatch

measures of ejaculatory latency. These measures are all relatively new and were developed primarily for use as research tools. Some have shown good psychometric properties and are potentially valuable adjuncts for clinical screening and assessment. Stopwatch measures of ejaculatory latency, in contrast, have been used extensively in research settings – both clinical trials and observational studies – but have typically not been recommended for use in clinical settings.

Several PE measures have been described in the literature⁹²⁻⁹⁸, although only a small number have undergone extensive psychometric testing and validation. Five validated questionnaires have been developed and published to date. Currently, there are two questionnaires that have extensive databases and meet most of the criteria for test development and validation: The Premature Ejaculation Profile (PEP) and the Index of Premature Ejaculation (IPE)^{92, 94}. A third brief diagnostic measure (PEDT) has also been developed, has a modest database and is available for clinical use⁹⁶. Two other measures (Arabic, Chinese PE Questionnaires) have minimal validation or clinical trial data available. These latter measures are not recommended for clinical use.

Premature Ejaculation Profile (PEP)

A four-item, self-report measure of premature ejaculation has been developed by Patrick et al⁹⁴. The PEP is comprised of single-item constructs of: (1) perceived control over ejaculation; (2) satisfaction with sexual intercourse; (3) personal distress related to ejaculation, (4) interpersonal difficulty related to ejaculation and (5) an index or total score. Each of the four individual items is assessed on a 5-point scale, which are averaged to provide an index PE score. The measure has been used in observational studies and clinical trials of premature ejaculation⁸⁴. It has also been recommended for clinical use in evaluating the subjective components of the disorder. Validation studies have been performed in comparison to stopwatch measures of intra-vaginal latency and other PRO measures of sexual function and distress^{84, 94}. The scale has adequate test-retest reliability (total scale = 0.80) and moderate to strong correlations with stopwatch measured IELT. A major limitation of the scale is the lack of validated cutoff scores, which make it less suitable for use as a diagnostic or clinical screening tool. On the positive side, it is very brief and easy to administer and may be valuable for use in a clinical setting as a measure of treatment responsiveness.

Index of Premature Ejaculation (IPE)

The Index of Premature Ejaculation (IPE) was developed by Althof et al⁹². It is a 10 item self-administered questionnaire designed to evaluate sexual satisfaction, control and distress in men with premature ejaculation. It was developed using four stages: item pool development, initial psychometric analyses, patient interviews, and final psychometric analyses. The IPE contains three factor analytically derived domains: control, sexual satisfaction and distress. All three domains have shown adequate internal consistency and reliability, as well as known groups validity in comparing men with and without PE. Convergent validity against IELT was also strong for all three domains (control (r=0.75); sexual satisfaction (r=0.60) and distress (r=0.68)).

The IPE has the advantages of also being relatively brief and easy-to-administer, although the measure is not as brief as the PEP above. It also assesses clinically relevant domains and

has adequate known groups validity data. However, similar to the PEP, it lacks norms and diagnostic cutoffs, and has limited value as a diagnostic or screening measure for PE.

Premature Ejaculation Diagnostic Tool (PEDT)

The previous two measures (PEP, IPE) are available for use as treatment change or outcome measures of PE treatment. The PEDT, in contrast, was developed specifically for use as a screening questionnaire^{95, 96, 98}. This questionnaire is a brief, 5-item measure used to screen men for potential presence of PE based on DSM-IV-TR criteria of lack of control, frequency, minimal stimulation, distress and interpersonal difficulty.

The measure has adequate internal consistency (Cronbach alpha = 0.71) and reliability (ICC = 0.73). Importantly known-groups validity is strong when comparing men with time-defined PE (IELTs ≤ 2 minutes in 70% of coital attempts) and those with self-reported no PE. A score of ≥ 11 would suggest presence of PE and men scoring 9 or 10 are likely to have PE and are recommended for further assessment. This scoring system has shown good convergent validity with clinical expert diagnosis of PE (k-statistic=0.80). The scale has also been translated into multiple languages.

Depending on the specific need, the PEP or IPE are currently the preferred questionnaire measures for assessing PE, particularly when monitoring responsiveness to treatment. Overall, these measures may serve as useful adjuncts, but should not substitute for a detailed sexual history performed by a qualified clinician. (LOE 2b)

The IPE, PEP and PEDT questionnaires can be found in Appendix 1

Treatment

Pharmacological Treatment

The use of anaesthetics to diminish the sensitivity of the glans penis is probably the oldest known form of treating premature ejaculation⁴⁵. The introduction of the selective serotonin reuptake inhibitors (SSRIs), paroxetine, sertraline, fluoxetine, citalopram and the tricyclic antidepressant (TCA) clomipramine has revolutionized the treatment of PE. These drugs block axonal re-uptake of serotonin from the synaptic cleft of central serotonergic neurons by 5-HT transporters, resulting in enhanced 5-HT neurotransmission and stimulation of post-synaptic membrane 5-HT autoreceptors.

Treatment with Selective Serotonin Reuptake Inhibitors (SSRIs) and Tricyclic Antidepressants (TCAs)

Premature ejaculation can be treated with on-demand SSRIs such as dapoxetine or off-label clomipramine, paroxetine, sertraline and fluoxetine, or with daily dosing of off-label paroxetine, clomipramine, sertraline, fluoxetine or citalopram.

Dapoxetine

Dapoxetine has received approval for the treatment of PE in Austria, Germany, Italy, Finland, Mexico, New Zealand, Portugal, South Korea, Spain, and Sweden. It is a rapid acting and short half-life SSRI with a pharmacokinetic profile suggesting a role as an on-demand treatment for PE^{19, 99-101}. No drug-drug interactions associated with dapoxetine, including phosphodiesterase inhibitor drugs, have been reported¹⁰². In RCTs, dapoxetine 30 mg or 60 mg taken 1-2 hr before intercourse is more effective than placebo from the first dose, resulting in a 2.5 and 3.0 fold increases in IELT, increased ejaculatory control, decreased distress, and increased satisfaction. Dapoxetine was comparably effective both in men with lifelong and acquired PE¹⁰³. Treatment related side effects were uncommon, dose dependent and included nausea, diarrhea, headache, and dizziness. They were responsible for study discontinuation in 4% (30mg) and 10% (60mg) of subjects. There was no indication of an increased risk of suicidal ideation or suicide attempts and little indication of withdrawal symptoms with abrupt dapoxetine cessation¹⁰⁴.

There is level 1a evidence to support the efficacy and safety of on-demand dosing of dapoxetine for the treatment of lifelong and acquired PE. (LOE 1a)

Off-label selective serotonin reuptake inhibitors (SSRIs) and Tricyclic Antidepressants (TCAs)

Daily treatment with off-label paroxetine 10-40 mg, clomipramine 12.5-50 mg, sertraline 50-200 mg, fluoxetine 20-40 mg, and citalopram 20-40 mg is usually effective in delaying ejaculation¹⁰⁵⁻¹¹⁰. A meta-analysis of published data suggests that paroxetine exerts the strongest ejaculation delay, increasing IELT approximately 8.8 fold over baseline²⁰.

Ejaculation delay usually occurs within 5-10 days of starting treatment, but the full therapeutic effect may require 2-3 weeks of treatment and is usually sustained during long-term use¹⁸. Adverse effects are usually minor, start in the first week of treatment and may gradually disappear within 2-3 weeks; they include fatigue, yawning, mild nausea, diarrhea or perspiration. Hypoactive desire and ED are infrequently reported and appear to have a lower incidence in non-depressed PE men compared to depressed men treated with SSRIs¹¹¹. Neurocognitive adverse effects include significant agitation and hypomania in a small number of patients, and treatment with SSRIs should be avoided in men with a history of bipolar depression¹¹².

Systematic analysis of RCTs in patients with depressive and/or anxiety disorders indicate a small increase in the risk of suicidal ideation or suicide attempts in youth but not adults¹¹³⁻¹¹⁵. In contrast, such risk of suicidal ideation has not been found in trials with SSRIs in non-depressed men with PE. Still caution is suggested in prescribing SSRIs to young adolescents with PE aged 18 years or less, and to men with PE and a comorbid depressive disorder, particularly when associated with suicidal ideation¹¹⁶. Patients should be advised to avoid sudden cessation or rapid dose reduction of daily dosed SSRIs which may be associated with a SSRI withdrawal syndrome¹¹⁷.

On-demand administration of clomipramine, paroxetine, sertraline and fluoxetine 3-6 hours before intercourse is modestly efficacious and well tolerated but is associated with substantially less ejaculatory delay than daily treatment in most studies¹¹⁸⁻¹²¹. On-demand treatment may be combined with either an initial trial of daily treatment or concomitant low dose daily treatment¹¹⁹.

Patients are reluctant to begin off-label treatment of PE with SSRIs. Salonia et al reported that 30% of patients refused to begin treatment (paroxetine, 10mg daily for 21 days followed by 20 mg as needed) and another 30% of those that began treatment discontinued it. Reasons given included: not wanting to take an antidepressant, treatment effects below expectations, temporary loss of interest in sex because of relationship issues and side effects¹²².

There is Level 1a evidence to support the efficacy and safety of off-label daily dosing of the SSRIs paroxetine, sertraline, citalopram, fluoxetine, and the serotonergic tricyclic, clomipramine, and off-label on-demand dosing of clomipramine, paroxetine, and sertraline for the treatment of lifelong and acquired PE. (LOE 1a)

The decision to treat PE with either on-demand dosing of dapoxetine (where available) or daily dosing of off-label SSRIs should be based upon the treating physician's assessment of individual patient requirements. Although many men with PE who engage in sexual intercourse infrequently may prefer on-demand treatment, many men in established relationships may prefer the convenience of daily medication. Well-designed preference trials will provide additional insight into the role of on-demand dosing.

In some countries off-label prescribing may present difficulties for the physician as the regulatory authorities strongly advise against prescribing for indications in which a medication is not licensed/approved. Obviously this complicates treatment in countries where there is no approved medication and the regulatory authorities advise against off-label prescription.

Topical Anesthetics

The use of topical local anesthetics such as lidocaine and/or prilocaine as a cream, gel, or spray is well established and is moderately effective in delaying ejaculation^{21, 123, 124}. Dinsmore et al reported on the use of PSD502 a lidocaine-prilocaine spray currently in clinical trials which is applied to the penis at least 5 minutes before intercourse. The treated group reported a 6.3 fold increase in IELT and associated improvements in PRO measures of control and sexual satisfaction²¹. There were minimal reports of hypoesthesias and transfer to the partner due to the unique formulation of the compound. Other topical anesthetics are associated with significant penile hypo-anesthesia and possible transvaginal absorption, resulting in vaginal numbness and resultant female anorgasmia unless a condom is used.

There is level 1b evidence to support the efficacy and safety of off-label on-demand label topical anaesthetics in the treatment of lifelong PE. (LOE 1b)

Phosphodiesterase Type 5 inhibitors

Phosphodiesterase type-5 inhibitors (PDE5i), sildenafil, tadalafil, and vardenafil, are effective treatments for ED. Several authors have reported using PDE5i's alone or in combination with SSRIs as a treatment for PE¹²⁵⁻¹²⁷. Although a review of 14 studies on the PDE5i drug treatment of PE has failed to provide robust empirical evidence to support a role of PDE5i in the treatment of PE with the exception of men with PE and co-morbid ED¹²⁸, recent well designed studies do support a potential role for these agents suggesting a need for further evidence based research¹²⁵.

There is level 4d evidence to support the efficacy and safety of off-label on-demand or daily dosing of PDE5i's in the treatment of lifelong PE in men with normal erectile function. Treatment of lifelong PE with PDE5i's in men with normal erectile function is not recommended and further evidence-based research is encouraged to understand conflicting data.

Other Pharmacological Treatments

Treatment with daily α 1-adrenoceptor antagonists, on-demand tramadol or intracavernous injection of vasoactive drugs has been reported in the literature¹²⁹⁻¹³¹. Salem et al reported that 25mg of tramadol, as needed, increased IELT from 1.17 minutes at baseline to 7.37 minutes after treatment¹³². Similarly, Safarinejad and Hosseini reported that 50mg of tramadol, as needed, resulted in IELT increases from 19 seconds at baseline to 243 seconds at the end of treatment. Twenty-eight percent of the tramadol group versus 15% of placebo patients reported treatment related adverse events including nausea, vomiting and dizziness¹³¹.

There is level 2d evidence to support the efficacy and safety of these treatments and their use as a treatment for PE cannot be recommended.

Table 4 is a summary of recommended pharmacological treatments for premature ejaculation

Surgery

Several authors have reported the use of surgically induced penile hypo-anesthesia via selective dorsal nerve neurotomy or hyaluronic acid gel glans penis augmentation in the treatment of lifelong PE refractory to behavioral and/or pharmacological treatment^{133, 134}. The role of surgery in the management of PE remains unclear until the results of further studies have been reported.

There is level 4 evidence i.e. no evidence, to suggest that selective dorsal nerve neurotomy or hyaluronic acid gel glans penis augmentation are effective treatments for PE. Surgery may be associated with permanent loss of sexual function and is currently not recommended in the management of PE.

Psychological/Behavioural, Combined Medical and Psychological and Educational Interventions

A wide range of psychological interventions have been developed for the treatment of PE. The majority of the psychotherapy treatment outcome studies are uncontrolled, unblinded

trials; none meet the requirements for high level evidence-based studies. The literature is comprised of reports on small to moderately sized cohorts of participants who received different forms of psychological interventions with limited or no follow-up. In most studies, active treatment was not compared to placebo, control or wait list groups¹³⁵. The most frequently used behavioral treatments are the squeeze or stop-start techniques^{33, 136}. Both of these therapies were designed to educate men to recognize when they are approaching the point of ejaculatory inevitability, and then either himself (if he is masturbating alone) or his partner (if she is masturbating him) stop the stimulation process until his sexual arousal subsides (generally about 30 seconds). Stimulation then resumes, until this point is reached again. This sequence of events is repeated 4-5 times after which the man ejaculates. A variation on this technique is used during intercourse, once the man has developed ejaculatory control during masturbation. This process gradually leads to an increase in the time to ejaculatory inevitability, teaches him to become more aware of his sexual excitement, and to control, his level of arousal. Further, this intervention is likely to lead to improvement in the man's confidence and self-esteem, although there are no controlled studies to support this claim.

Older uncontrolled studies on the squeeze technique report a failure rate of 2.2% immediately after therapy, and 2.7% at the 5-year follow-up³³. These results have not been replicated; other studies have found success rates of between 60% and 90%¹³⁷. In a recent study Carufel and Trudel demonstrated an eight fold increase in IELT among men treated with the above behavioral techniques compared to a wait list control condition¹⁷.

Psychological interventions are designed to achieve more than simply increasing the IELT. Targeted factors focus on the man, his partner and their relationship. In particular, they include 1) an increase in man's confidence in this sexual performance, as well as his overall self-confidence; 2) lowered performance anxiety; 3) an increase in communication with the partner; and 4) a resolution of interpersonal problems that may have precipitated or maintained the PE.

There is level 2b evidence regarding the efficacy of psychological/behavioral interventions in the treatment of PE.

Men with Natural Variable PE (irregular and inconsistent rapid ejaculation with a diminished sense of subjective control of ejaculation) should be educated and reassured. Men with Premature Like Ejaculatory Dysfunction (those whose IELT are within the normal range but who are preoccupied by their ejaculatory control) may require a referral for psychotherapy. More research is necessary to better define the efficacy of reassurance, education and psychotherapy with these provisional subtypes.

Primary care physicians, urologists and mental health professionals have differing levels of interest and training in treating PE. In general all clinicians should be able to diagnose, offer support and prescribe behavioral exercises. If the situation is complex and/or patients are not responsive the initial intervention clinicians should consider referring to a sexual health specialist.

Importance of Partners

Inclusion of the partner in the treatment process is an important but not a mandatory ingredient for treatment success¹³⁸. Some patients may not understand why the clinician wishes to include the partner and some partners may be reluctant to join the patient in treatment. However, if partners are not involved in treatment, they may be resistant to changing the sexual interaction. A cooperative partner can enhance the man's self-confidence, skills, self-esteem, sense of masculinity, and more generally assist the man to develop ejaculatory control. This is, in turn, likely to lead to an improvement in the couple's sexual relationship, as well as the broader aspects of their relationship. There are no controlled studies on the impact of involving partners in treating PE. However, a review of treatment studies for ED demonstrated the important role of including a focus on interpersonal factors on treatment success¹³⁹.

Benefits of Combination Medical and Psychological Therapy

There are three studies reporting on combined pharmacological and behavioral treatment for PE¹⁴⁰⁻¹⁴² and one study reporting on the consecutive treatment with pharmacotherapy followed by behavior therapy¹⁴³. Each study reported on a different medication- sildenafil, citalopram, clomipramine or paroxetine (in the consecutive study). Pharmacotherapy was given in conjunction with a behavioral treatment and compared to pharmacotherapy alone. **In all three studies combination therapy was superior to pharmacotherapy alone on either IELT and/or the Chinese Index of Premature Ejaculation.**

For ED, combined treatments have also been found to be more effective than either medical or psychological treatments alone¹⁴⁴⁻¹⁴⁶. Factors that are not addressed by medical treatments that can be treated with a psychological approach include: 1) patient factors (e.g., performance anxiety, self-confidence); 2) partner factors (e.g., partner sexual dysfunction); 3) relationship factors (e.g., conflict, lack of communication); 4) sexual factors in the relationship (e.g., sexual scripts, sexual satisfaction); and 5) contextual factors (e.g., life stressors).

Combining a medical and psychological approach may be especially useful in men with acquired premature ejaculation where there is a clear psychosocial precipitant or lifelong cases where the individual or couple's responses to PE are likely to interfere in the medical treatment and ultimate success of therapy. Similarly, in men with PE and co-morbid ED combination therapy may also be helpful to manage the psychosocial aspects of these sexual dysfunctions. (LOE 2a)

Role of Education and Coaching

Education on PE (or coaching) may also be useful in the treatment of this condition¹⁴⁷⁻¹⁵⁰. Such techniques may treat those aspects of PE that are not treated with medication. They include providing information on the prevalence of PE, and general population IELT to dispel myths about PE, information on enjoyable sexual activities to extend the man and his partner's sexual repertoire, as well as strategies to address avoidance of sexual activity or unwillingness to discuss sex with his partner. These **educational strategies are designed**

to give the man the confidence to try the medical intervention, reduce performance anxiety, and modify his maladaptive sexual scripts. (LOE 5d)

Life-long PE

Since life-long PE is likely to have an organic etiology, a medical intervention with basic psycho-education is initially recommended^{36, 151}. However, if the PE has resulted in psychological and relationship concerns, graded levels of patient and couple counselling, guidance and/or relationship therapy may be a useful adjunct to the medical intervention. (LOE 1a)

Acquired PE

It is recommended that HCPs utilize a combination medical and psychological approach where feasible. Men desire an immediate effect from therapy; therefore medical therapy and amelioration of associated disease factors such as ED will be extremely helpful.

Additionally, education on the nature of PE, helping men improve ejaculatory control with behavioral exercises, addressing restricted/narrow sexual behavioral patterns and resolving interpersonal issues are likely to be of significant help to men with acquired PE. Once the man's self confidence and sense of control have improved, it may then be possible to reduce or discontinue the medical intervention¹⁸. (LOE 5d)

Special Patient Populations

Premature Ejaculation and Co-morbid ED

Recent data demonstrate that as many as 30-50% of subjects with ED also experience PE^{9, 10}. Subjects with ED may either require higher levels of manual stimulation to achieve an erection or intentionally "rush" intercourse to prevent early detumescence of a partial erection, resulting in ejaculation with a brief latency. This may be compounded by the presence of high levels of performance anxiety related to their ED which serves to only worsen their prematurity.

There is evidence to suggest that a PDE5i alone or in combination with a SSRI may have a role in the management of acquired PE in men with co-morbid ED¹⁵²⁻¹⁵⁴. The high correlation between PDE5i improved erectile function and increased IELTs indicates that the PDE5i-related reduced PE severity is due to improved erectile function^{153, 154}

While men with mild ED and PE do benefit from treatment with SSRIs, compared to men with no ED, their response to treatment is diminished¹⁵². Additionally, men with lifelong PE and mild ED were less responsive to treatment than men with acquired PE and mild ED¹⁰³.

There is Level 1A evidence to support the treatment of PE and co-morbid ED with ED pharmacotherapy. There is Level 3C evidence to support the treatment of PE

and co-morbid ED with ED pharmacotherapy in combination with PE pharmacotherapy. Further evidence-based research is encouraged

Premature Ejaculation and Hyperthyroidism

There is evidence to indicate a link between depression, serotonin and thyroid hormones^{155, 156}. While many patients with hypothyroidism experience erectile dysfunction, few experience PE. Waldinger et al studied 620 consecutive men with lifelong PE without erectile dysfunction and found no relationship between hyper- or hypothyroidism and lifelong PE⁷³. However, Carini et al studied 48 men with hyper and hypothyroidism and found that 50% of hyperthyroid men had acquired PE. Treatment of the hyperthyroidism with thyroid hormone normalization using anti-thyroid drugs, radioactive iodine or thyroidectomy resulted in normalization of ejaculatory function in 35% of these men⁵⁷. This finding requires further replication and at this point the Committee does not recommend routine TSH screening in men with acquired PE.

Premature Ejaculation and Chronic Prostatitis

Although antibiotic treatment of chronic prostatitis improves LUTS, there is little published data to suggest a parallel improvement in PE and other sexual dysfunction symptoms¹⁵⁷⁻¹⁵⁹. Although physical and microbiological examination of the prostate in men with painful ejaculation or LUTS is mandatory, there is insufficient evidence to support routine screening of men with PE for chronic prostatitis.

Outcome

Across all treatments, including daily and as-needed dosing schedules patient's IELT may be expected to increase between 3 to 8 fold compared to their baseline IELT. Similarly, patient's perceived control (self efficacy) of their ejaculatory timing was significantly improved with treatment. From the patient's perspective, perceived ejaculatory control was a more important outcome variable than IELT alone¹⁶⁰.

Treatment outcome can be addressed in one simple, brief and validated question known as the Clinical Global Index of Change (CGIC)¹⁶¹. It asks patients, "Compared to before starting treatment, would you describe your premature ejaculation problem as: much worse, worse, slightly worse, no change, slightly better, better, or much better?" In the original research study the phrase "before starting treatment" was actually, "before the start of the study. Further research should document the validity of this minor wording change. (LOE 1b)

In randomized, placebo-controlled, double-blind trials of dapoxetine for each category of improvement on the CGIC, the magnitude of IELT prolongation increased in concert with the CGIC rating. Additionally, significant positive correlations were noted between CGIC ratings and ejaculatory control and sexual satisfaction and inverse correlation were noted with sexual distress. Control accounted for the greatest proportion of the variance of the CGIC.

Role of the Primary Care Clinician (PCP)

Primary care providers are usually the first line of contact for a patient with the health care system including the diagnosis and treatment of sexual problems. This role includes 1) initial recognition and evaluation of any undiagnosed sign, symptom, or health concern (the "undifferentiated" patient); 2) health promotion including disease prevention, health maintenance, counseling, patient education, chronic illness management, and patient advocacy; and 3) coordination of care promoting effective communication with patients and encouraging the patient to be a partner in health¹⁶². This model of care is not limited by problem origin, organ system, or diagnosis.

Primary care clinicians are the ideal group to assist the patient with sexual difficulties for several reasons including: 1) the value of the longitudinal and personal relationships with patients in discussing and resolving sexual problems; 2) the multifactorial issues around sexual problems that can be appropriately evaluated by a generalist clinician; and; 3) the long-term follow-up routine in primary care is well-suited to being certain that a sexual dysfunction is resolved. The main obligation of the PCP is to recognize PE and make the patient feel comfortable about getting help. Initial work-up and treatment can be planned by the PCP who has good communication skills about sexual activity and who is knowledgeable about first-line treatments. Urologists can be helpful in difficult or complex situations or when the patient presents with a genital anatomic problem or when a patient presents with ED or painful intercourse due to phimosis. A mental health professional with experience in sexual problems can help working collaboratively with the clinician to increase the chances of therapeutic success by: 1) resolving the sexual difficulty; 2) teasing out important history; 3) educating the patient and partner; 4) suggesting sexual enhancement techniques; and 5) helping the couple resolve individual as well as relationship problems.

Help from a sexual health specialist may be appropriate at varying intervals when managing a man with premature ejaculation. The major factors in determining when a consultation is needed include 1) the primary care clinician's comfort in discussing and managing treatment options; 2) the depth of the psychosocial and sexual issues involved; and 3) the success or failure of the initial intervention efforts. Specific instances in which subspecialty assistance is often useful include 1) treatment failure; 2) anatomic or complex hormonal issues; 3) complex issues around sex and/or partnerships; 4) severe psychological problems; or 5) anytime the treating physician feels that help is needed

The management algorithm "**ALLOW**" a outgrowth of the PLISSIT model(permission, limited information, specific suggestions and intensive therapy)¹⁶³ can help PCPs talk with their patients about sexual health problems and refer patients when appropriate¹⁶⁴. The physician begins by "**A**sking" about a patient's sexual life and "**L**egitimizes" the importance and potential impact of sexual problems to the patient. The physician then considers his/her "**L**imitations" with regard to managing the sexual dysfunction, and may refer the patient to a sexual health specialist for further evaluation and management. Conversely, if the physician feels comfortable managing the patient's issue(s), he/she then "**O**pens: up the issue(s) for further discussion, and the physician and the patient "**W**ork together to develop a management plan." This methodical approach to "allowing" the patient to discuss sexual issues can be accomplished in as brief or as long a time as the physician has available.

When a clinical situation is encountered in which the primary care clinician needs assistance in management, two options are available. Patients can be sent to a sexual health specialist for a “consultation.” This is a request to answer a specific question with the intention that management will be advanced under the primary care clinician considering the added input. The second option is “referring” the patient to another clinician, usually a sexual health specialist in the field involved, to move further with management. In these situations, a summary of the reason for referral, appropriate history, and related diagnostic studies should accompany the patient. The sexual health specialist will then appropriately advance the management keeping the primary care clinician advised of progress. In a “referral” the care for the referred patient is usually transferred to the sexual health specialist. Communication can be enhanced by the primary care clinician by specifically informing the sexual health specialist about the level of involvement requested¹⁶⁵.

Co-management by the less experienced primary care clinician along with an appropriate sexual health specialist is an excellent way to manage the patient’s issues and to increase the primary care clinician’s understanding of PE. Good communication between clinicians can improve patient outcome and understanding of planned treatments and monitoring measures. Co-management can be optimized by a clear understanding between the primary care clinician and the sexual health specialist about who will do what. This kind of communication can be ensured by creating “referral agreements” in which a specific referral guideline is used and the responsibilities and activities of each clinician are clearly spelled out¹⁶⁶.

Conclusion

These guidelines have been carefully promulgated by an interdisciplinary international panel of recognized experts in the area of PE with the goal of constructing clearly worded, practical, evidenced-based recommendations for the diagnosis and treatment of PE for family practice clinicians as well as sexual medicine experts. Recognizing that all evidence is not created equal the Committee reviewed and debated the research literature to arrive at its grading of their recommendations according to the Oxford Centre of Evidence-Based Medicine rankings. Table 1 lists all the relevant recommendation of the PE Guidelines Committee. These Guidelines affirm the ISSM definition of premature ejaculation and suggest that the prevalence is considerably lower than previously thought. Evidence-based data regarding biological and psychological etiology of PE are presented, as is population-based statistics on normal ejaculatory latency. Brief assessment procedures are delineated and validated diagnostic and treatment questionnaires are reviewed. Finally, the best practices treatment recommendations are presented that aim to guide clinicians, both familiar and unfamiliar with premature ejaculation, in facilitating treatment of their patients.

Development of guidelines is an evolutionary process that continually reviews data and incorporates the best new research. We expect that ongoing research will lead to a more complete understanding of the pathophysiology as well as new efficacious and safe treatments for this sexual dysfunction. Therefore, it is strongly recommended that these guidelines be re-evaluated and updated by the ISSM every 4 years.

Finally, it is important to keep in mind that premature ejaculation causes significant personal and interpersonal distress to the man, his partner and the couple. We are hopeful that these guidelines will assist clinicians in accurately diagnosing and managing their patients who present with complaints of PE.

References

- 1 Rowland D, Mc Mahon C, Abdo C, Chen J, Jannini E, Waldinger M, et al. Disorders of orgasm and ejaculation in men. *Journal of Sexual Medicine*. 2010;**7**: 1668-86.
- 2 Borgdorff AJ, Bernabe J, Denys P, Alexandre L, Giuliano F. Ejaculation elicited by microstimulation of lumbar spinothalamic neurons. *European Urology*. 2008;**54**: 449-56.
- 3 Clement P, Peeters M, Bernabe J, Denys P, Alexandre L, Giuliano F. Brain oxytocin receptors mediate ejaculation elicited by 7-hydroxy-2-(di-N-propylamino) tetralin (7-OH-DPAT) in anaesthetized rats. *British Journal of Pharmacology*. 2008;**154**: 1150-9.
- 4 Peeters M, Giuliano F. Central neurophysiology and dopaminergic control of ejaculation. *Neuroscience Biobehavioral Review*. 2008;**32**: 438-53.
- 5 Sun XQ, Xu C, Leclerc P, Benoit G, Giuliano F, Droupy S. Spinal neurons involved in the control of the seminal vesicles: a transsynaptic labeling study using pseudorabies virus in rats. *Neuroscience*. 2009;**158**: 786-97.
- 6 Truitt W, Coolen L. Identification of a potential ejaculation generator in the spinal cord. *Science*. 2002;**297**: 1566-69.
- 7 Coolen L, Allard J, Truitt W, McKenna K. Central regulation of ejaculation. *Physiology and Behavior*. 2004;**83**: 203-15.
- 8 Coolen L. Neural control of ejaculation. *Journal of Comparative Neurology*. 2005;**493**: 39-45.
- 9 Laumann EO, Nicolosi A, Glasser DB, Paik A, Gingell C, Moreira E, et al. Sexual problems among women and men aged 40-80 y: prevalence and correlates identified in the Global Study of Sexual Attitudes and Behaviors. *International Journal of Impotence Research*. 2005;**17**: 39-57.
- 10 Porst H, Montorsi F, Rosen RC, Gaynor L, Grupe S, Alexander J. The premature ejaculation prevalence and attitudes (PEPA) survey: Prevalence, comorbidities and professional help-seeking. *European Urology*. 2007;**51**: 816-24.
- 11 Waldinger M, Quinn P, Dilleen M, Mundayat R, Schweitzer D, Boolell M. A multinational population survey of intravaginal ejaculation latency time. *Journal of Sexual Medicine*. 2005;**2**: 292-97.
- 12 Nicolosi A, Laumann E, Glaser D, Moreira E, Paik A, Gingell C. Global study of sexual attitudes and behaviors investigator's group. Sexual behavior and sexual dysfunctions after age 40: the global study of sexual attitudes and behaviors. *Urology*. 2004;**64**: 991-97.
- 13 Lindau S, Schumm P, Laumann E, Levinson W, O'Muirheartaigh C, Waite L. A Study of Sexuality and Health among Older Adults in the United States. *New England Journal of Medicine*. 2007;**357**: 762-64.
- 14 McMahon CG, Althof SE, Waldinger MD, Porst H, Dean J, Sharlip ID, et al. An evidence-based definition of lifelong premature ejaculation: report of the International Society for Sexual Medicine (ISSM) ad hoc committee for the definition of premature ejaculation. *Journal of Sexual Medicine*. 2008;**5**: 1590-606.
- 15 Rosen R, Althof S. Impact of premature ejaculation: The psychological quality of life and sexual relationship consequences. *Journal of Sexual Medicine*. 2008;**5**: 1296-307.
- 16 Althof S, Symonds T. Patient reported outcomes used in the assessment of premature ejaculation. In: Seftel A, ed. *Urologic Clinics of North America*; 2007:581-89.
- 17 De Carufel F, Trudel G. Effects of a new functional sexological treatment for premature ejaculation. *Journal of Sex and Marital Therapy*. 2006;**32**: 97-114.
- 18 McMahon CG. Long term results of treatment of premature ejaculation with selective serotonin reuptake inhibitors. *International Journal of Impotence Research*. 2002;**14**: S19.
- 19 Pryor JL, Althof SE, Steidle C, Rosen RC, Hellstrom WJ, Shabsigh R, et al. Efficacy and tolerability of dapoxetine in treatment of premature ejaculation: an integrated analysis of two double-blind, randomised controlled trials. *Lancet*. 2006;**368**: 929-37.
- 20 Waldinger MD, Zwinderman AH, Schweitzer DH, Olivier B. Relevance of methodological design for the interpretation of efficacy of drug treatment of premature ejaculation: a systematic review and meta-analysis. *International Journal of Impotence Research*. 2004;**16**: 369-81.
- 21 Dinsmore WW, Hackett G, Goldmeier D, Waldinger M, Dean J, Wright P, et al. Topical eutectic mixture for premature ejaculation (TEMPE): a novel aerosol-delivery form of lidocaine-prilocaine for treating premature ejaculation. *British Journal of Urology, International*. 2007;**99**: 369-75.
- 22 American Psychological Association. Criteria for practice guideline development and evaluation. *American Psychologist*. 2002;**57**: 1048-51.
- 23 Lin K, Slawson D. Identifying and using good practice guidelines. *American Family Physician*. 2009;**80**: 67-69.

- 24 Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is
and what it isn't. *British Medical Journal*. 1996;**312**: 71-2.
- 25 Ebell M, Siwek J, Weiss B, Woolf S, Susman J, Ewigman B, et al. Strength of recommendation taxonomy
(SORT): A patient-centered approach to grading evidence in the medical literature. *Journal of the
American Board of Family Practice*. 2004;**17**: 59-67.
- 26 Hatzimouratidis K, Amar E, Eardley I, Giuliano F, Hatzichristou D, Montorsi F, et al. Guidelines on Male
Sexual Dysfunction: Erectile Dysfunction and Premature Ejaculation. *European Urology*. 2010: epub
- 27 Ghanem H, El-Meliegy A, El Sakka A, Jad A. Practice guidelines for the Pan-Arab Society of Sexual
Medicine. 2005.
- 28 Montague D, Jarow J, Broderick G, Dmochowski R, Heaton J, Lue T. AUA guideline on the
pharmacologic management of premature ejaculation. *Journal of Urology*. 2004;**172**: 290-94.
- 29 Colpi G, Weidner W, Jungwirth A, Pomerol J, Papp G, Hargreave T, et al. EAU guidelines on ejaculatory
dysfunction. *European Urology*. 2004;**46**: 555-58.
- 30 Oxford Centre for Evidence Based Medicine Levels of Evidence www.cebm.net. 2001.
- 31 Althof SE, Abdo CH, Dean J, Hackett G, McCabe M, McMahon CG, et al. International Society for Sexual
Medicine's guidelines for the diagnosis and treatment of premature ejaculation. *J Sex Med*.**7**: 2947-69.
- 32 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, 4th Edition, Text
Revision*. Washington, DC: American Psychiatric Association; 2000.
- 33 Masters W, Johnson V. *Human Sexual Inadequacy*. Boston: Little, Brown; 1970.
- 34 Metz M, McCarthy B. *Coping with premature ejaculation: How to overcome PE, please your partner &
have great sex*. Oakland: New Harbinger Publications; 2003.
- 35 World Health Organization. International Classification of Diseases and Related Health Problems. 10th
edition Geneva; 1994.
- 36 Waldinger M. Premature ejaculation: Different pathophysiologies and etiologies determine its treatment.
Journal of Sex and Marital Therapy. 2008;**34**: 1-13.
- 37 Pagani E, Rodrigues O, Torselli M, Genari D. Characterization of 305 Men With Complaints of Premature
Ejaculation. *International Journal of Impotence Research*. 1996;**8**: 172.
- 38 Serefoglu EC, Cimen HI, Atmaca AF, Balbay MD. The Distribution of Patients Who Seek Treatment for
the Complaint of Ejaculating Prematurely According to the Four Premature Ejaculation Syndromes.
Journal of Sexual Medicine. 2010;**7**: 810-15.
- 39 Montorsi F. Prevalence of premature ejaculation: A global and regional perspective. *Journal of Sexual
Medicine*. 2005;**Suppl 2**.
- 40 Jannini E, Lenzi A. Epidemiology of premature ejaculation. *Current Opinion in Urology*. 2005;**15**: 399-
403.
- 41 Abdo C. Estudo da Vida Sexual do Brasileiro. São Paulo: Bregantini; 2004.
- 42 Brock G, Benard F, Casey R, Elliott S, Gajewski J, Lee J. Canadian male sexual health council survey to
assess prevalence and treatment of premature ejaculation in Canada *Journal of Sexual Medicine*.
2009;**6**: 2115-23.
- 43 Waldinger M, McIntosh J, Schweitzer DH. A five-nation survey to assess the distribution of the
intravaginal ejaculatory latency time among the general male population. *Journal of Sexual Medicine*.
2009;**6**: 2888-95.
- 44 Shindel A, Nelson C, Brandes S. Urologist practice patterns in the management of premature
ejaculation: a nationwide survey. *Journal of Sexual Medicine*. 2008;**5**: 199-205.
- 45 Schapiro B. Premature ejaculation, a review of 1130 cases. *Journal of Urology*. 1943;**50**: 374- 79.
- 46 Xin Z, Choi Y, Rha K. Somatosensory evoked potentials in patients with primary premature ejaculation
Journal of Urology. 1997;**158**: 451-55.
- 47 Fanciullacci F, Colpi GM, Beretta G, Zanollo A. Cortical evoked potentials in subjects with true premature
ejaculation. *Andrologia*. 1988;**20**: 326-30.
- 48 Waldinger M, Berendsen HH, Blok BF, Olivier B, Holstege G. Premature ejaculation and serotonergic
antidepressants-induced delayed ejaculation: the involvement of the serotonergic system. *Behavioral
Brain Research*. 1998;**92**: 111-18.
- 49 Giuliano F. 5-Hydroxytryptamine in premature ejaculation: opportunities for therapeutic intervention. *Trends in
Neuroscience*. 2007;**30**: 79-84.
- 50 Chia S. Management of premature ejaculation: A comparison of treatment outcome in patients with and
without erectile dysfunction. *International Journal of Andrology*. 2002;**25**: 301-05.

- 51 Screponi E, Carosa E, Di Stasi SM, Pepe M, Carruba G, Jannini EA. Prevalence of chronic prostatitis in men with premature ejaculation. *Urology*. 2001;**58**: 198-202.
- 52 Xing JP, Fan JH, Wang MZ, Chen XF, Yang ZS. [Survey of the prevalence of chronic prostatitis in men with premature ejaculation]. *Zhonghua Nan Ke Xue*. 2003;**9**: 451-3.
- 53 O'Flynn R, Michael A. Reboxetine-induced spontaneous ejaculation. *British Journal of Psychiatry*. 2000;**177**: 567-8.
- 54 Adson D, Kotlyar M. Premature ejaculation associated with citalopram withdrawal. *Annals of Pharmacotherapy*. 2003;**37**: 1804-06.
- 55 Peugh J, Belenko S. Alcohol, drugs and sexual function: a review. *Journal of Psychoactive Drugs*. 2001;**33**: 223-32.
- 56 Gonen M, Kalkan M, Cenker A, Ozkardes H. Prevalence of premature ejaculation in Turkish men with chronic pelvic pain syndrome. *Journal of Andrology*. 2005;**26**: 601-3.
- 57 Carani C, Isidori AM, Granata A, Carosa E, Maggi M, Lenzi A, et al. Multicenter Study on the Prevalence of Sexual Symptoms in Male Hypo- and Hyperthyroid Patients. *Journal of Clinical Endocrinology and Metabolism*. 2005;**90**: 6472-79.
- 58 Corona G, Petrone L, Mannucci E, Jannini E, Mansani R, A M, et al. Psycho-biological correlates of rapid ejaculation in patients attending an andrologic unit for sexual dysfunctions. *European Urology*. 2004;**46**: 615-22.
- 59 Waldinger M, Schweitzer DH. The use of old and recent DSM definitions of premature ejaculation in observational studies: a contribution to the present debate for a new classification of PE in the DSM-V. *Journal of Sexual Medicine*. 2008;**5**: 1079-87.
- 60 Jern P, Santtila P, Alanko K, Harlaar N, Johansson A, von der Pahlen B, et al. Premature and delayed ejaculation: Genetic and environmental effects in a population-based sample of Finnish twins. *Journal of Sexual Medicine*. 2007;**4**: 1739-49.
- 61 Waldinger M. The neurobiological approach to premature ejaculation. *Journal of Urology*. 1998;**168**: 2359-67.
- 62 Borgdorff A, Rossler A, Clement P, Bernabe J, Alexandre L, Giuliano F. Differences in the spinal command of ejaculation in rapidly ejaculating rats. *Journal of Sexual Medicine*. 2009;**6**: 2197-205.
- 63 Waldinger M, Rietschel M, Nothen N, Hengeveld MW, Olivier B. Familial occurrence of primary premature ejaculation. *Psychiatric Genetics*. 1998;**8**: 37-40.
- 64 Janssen P, Bakker S, Rethelyi J, Zwinderman A, Touw D, Olivier B, et al. Serotonin transporter promoter region (5-HTTLPR) polymorphism is associated with the intravaginal ejaculation latency time in Dutch men with lifelong premature ejaculation. . *Journal of Sexual Medicine*. 2009;**6**: 276-84.
- 65 Ozbek E, Tasci A, Tugcu V, Ilbey Y, Simsek A, Ozcan L, et al. Possible association of the 5-HTTLPR serotonin transporter promoter gene polymorphism with PE in a Turkish population. *Asian Journal of Andrology*. 2009: 1-5.
- 66 Safarinejad M. Polymorphisms of the serotonin transporter gene and their relation to premature ejaculation in individuals from Iran. *Journal of Urology*. 2009;**181**: 2656-61.
- 67 Waldinger M, Janssen P, Schweitzer D. Hardy Weinberg equilibrium in genetic PE research remains critical to avoid misinterpretation. Author Reply. *Asian Journal of Andrology*. 2009;**11**: 524.
- 68 Waldinger MD, Janssen PK, Schweitzer DH. Re: Polymorphisms of the serotonin transporter gene and their relation to premature ejaculation in individuals from Iran. M. R. Safarinejad. *J Urol* 2009; 181: 2656-2661. *J Urol*. 2009;**182**: 2983; author reply 83-4.
- 69 Kulikov AV, Maksyutova AV, Ivanova EA, Khvorostov IB, Popova NK. The effect of thyroidectomy on the expression of the mRNA of 5-HT_{2A} serotonin receptors in the rat frontal cortex. *Dokl Biochem Biophys*. 2002;**383**: 116-8.
- 70 Kulikov AV, Zubkov EA. Chronic thyroxine treatment activates the 5-HT_{2A} serotonin receptor in the mouse brain. *Neurosci Lett*. 2007;**416**: 307-9.
- 71 Corona G, Petrone L, Mannucci E, Jannini EA, Mansani R, Magini A, et al. Psycho-biological correlates of rapid ejaculation in patients attending an andrologic unit for sexual dysfunctions. *Eur Urol*. 2004;**46**: 615-22.
- 72 Corona G, Jannini E, Lotti F, Boddi V, DeVita G, Forti G, et al. Premature and delayed ejaculation: two ends of a single continuum influenced by hormonal milieu. . *International Journal of Andrology*. 2010: epub ahead of print.

- 73 Waldinger M, Zwinderman A, Olivier B, Schweitzer D. Thyroid stimulating hormone assessments in a Dutch cohort of 620 men with lifelong premature ejaculation without erectile dysfunction. *Journal of Sexual Medicine*. 2005;**2**: 865-70.
- 74 Liang CZ, Zhang XJ, Hao ZY, Shi HQ, Wang KX. Prevalence of sexual dysfunction in Chinese men with chronic prostatitis. *British Journal of Urology, International*. 2004;**93**: 568-70.
- 75 Trinchieri A, Magri V, Cariani L, Bonamore R, Restelli A, Garlaschi MC, et al. Prevalence of sexual dysfunction in men with chronic prostatitis/chronic pelvic pain syndrome. *Archives of Italian Urology and Andrology*. 2007;**79**: 67-70.
- 76 Shamloul R, el-Nashaar A. Chronic prostatitis in premature ejaculation: a cohort study in 153 men. *Journal of Sexual Medicine*. 2006;**3**: 150-54.
- 77 Lotti F, Corona G, Mancini M, Biagini C, Colpi G, Innocenti S, et al. The association between varicocele, premature ejaculation and prostatitis symptoms: possible mechanisms. *Journal of Sexual Medicine*. 2009;**6**: 2878-87.
- 78 Althof S, Leiblum S, Chevret M, Hartmann U, Levine SB, McCabe M, et al. Psychological and Interpersonal Dimensions of Sexual Function and Dysfunction. In: Lue T, Basson R, Rosen R, Giuliano F, Khoury S, Montorsi F, eds. *Sexual Medicine: Sexual Dysfunctions in Men and Women*. Paris: Edition 2004; 2004.
- 79 McCabe M. The development and maintenance of sexual dysfunction: An explanation based on cognitive theory. *Sex and Marital Therapy*. 1991;**6**: 254-60.
- 80 Michetti P, Rossi R, Bonanno D, DeDominicis C, Lori F, Simonelli C. Dysregulation of emotions and premature ejaculation (PE): Alexithymia in 100 outpatients. *Journal of Sexual Medicine*. 2007;**17**: 18-23.
- 81 Symonds T, Roblin D, Hart K, Althof S. How does premature ejaculation effect a man's life. *Journal of Sex and Marital Therapy*. 2003;**29**: 361-70.
- 82 Rowland D, Patrick D, Rothman M, Gagnon D. The psychological burden of premature ejaculation. *Journal of Urology*. 2007;**177**: 1065-70.
- 83 Hobbs K, Symonds T, Abraham L, May K, Morris MF. Sexual dysfunction in partners of men with premature ejaculation. *Int J Impot Res*. 2008;**20**: 512-7.
- 84 Patrick DL, Althof SE, Pryor JL, Rosen R, Rowland DL, Ho KF, et al. Premature ejaculation: an observational study of men and their partners. *Journal of Sexual Medicine*. 2005;**2**: 358-67.
- 85 Schein M, Zyzanski SJ, Levine S, Medalie JH, Dickman RL, Alemagno SA. The frequency of sexual problems among family practice patients. *Journal of Family Practice Research*. 1988;**7**: 122-34.
- 86 Humphrey S, Nazareth I. GP's view on their management of sexual dysfunction. *Family Practice*. 2001;**18**: 516-18.
- 87 Mc Mahon C, Abdo C, Incrocci L, Perelman M, Rowland D, Stuckey B, et al. Disorders of orgasm and ejaculation in men. In: Lue T, Basson R, Rosen R, eds. *Sexual Medicine: Sexual Dysfunctions in Men and Women (2nd International Consultation on Sexual Dysfunctions)*. Paris: Health Publications; 2004:409-68.
- 88 Althof SE. Evidence based assessment of rapid ejaculation. *Int J Impot Res*. 1998;**10 Suppl 2**: S74-6; discussion S77-9.
- 89 Pryor JL, Broderick GA, Ho KF, Jamieson C, Gagnon D. Comparison of estimated versus measured intravaginal ejaculatory latency time in men with and without premature ejaculation. *Journal of Sexual Medicine*. 2005;**3**: 54.
- 90 Rosen R, McMahon C, Niederberger C, Broderick G, Jamieson C, Gagnon DD. Correlates to the clinical diagnosis of premature ejaculation: Results from a large observational study of men and their partners. *Journal of Urology*. 2007;**177**: 1059-64.
- 91 McMahon C. Clinical trial methodology in premature ejaculation observational, interventional and treatment preference studies- Part 1- Defining and selecting the study population. *Journal of Sexual Medicine*. 2008;**6**: 1805-16.
- 92 Althof S, Rosen R, Symonds T, Mundayat R, May K, Abraham L. Development and validation of a new questionnaire to assess sexual satisfaction, control and distress associated with premature ejaculation. *Journal of Sexual Medicine*. 2006;**3**: 465-75.
- 93 Arafa M, Shamloul R. Development and validation of the Arabic Index of Premature Ejaculation (AIPE). *Journal of Sexual Medicine*. 2007;**4**: 1750-56.

- 94 Patrick DL, Giuliano F, Ho KF, Gagnon DD, McNulty P, Rothman M. The Premature Ejaculation Profile: Validation of self-reported outcome measures for research and practice. *British Journal of Urology, International*. 2008;**103**: 358-67.
- 95 Symonds T, Perelman M, Althof S, Giuliano F, Martin M, Abraham L, et al. Further evidence of the reliability and validity of the premature ejaculation diagnostic tool. *International Journal of Impotence Research*. 2007;**19**: 521-25.
- 96 Symonds T, Perelman M, Althof S, Giuliano F, Martin M, May K, et al. Development and validation of a premature ejaculation diagnostic tool. *European Urology*. 2007;**52**: 565-73.
- 97 Yuan Y, Xin ZC, Jiang J, Guo Y, Liu W, Tian L, et al. Sexual function of premature ejaculation patients assayed with the Chinese Index of Premature Ejaculation. *Asian Journal of Andrology*. 2004;**6**: 121-26.
- 98 Serefoglu E, Cimen H, Ozdemir A, Symonds T, Berktaş M, Balbay M. Turkish validation of the Premature Ejaculation Diagnostic Tool and its association with intravaginal ejaculatory latency time. *International Journal of Impotence Research*. 2009;**21**: 139-44.
- 99 Buvat J, Tesfaye F, Rothman M, Rivas DA, Giuliano F. Dapoxetine for the treatment of premature ejaculation: results from a randomized, double-blind, placebo-controlled phase 3 trial in 22 countries. *European Urology*. 2009;**55**: 957-67.
- 100 Hellstrom WJ, Althof S, Gittelman M, Streidle C, Ho KF, Kell S, et al. Dapoxetine for the treatment of men with premature ejaculation (PE):dose-finding analysis. *Journal of Urology*. 2005;**173**: 238 -abstract 877.
- 101 McMahon C, Kim S, Park N, Chang C, Rivas D, Tesfaye F, et al. Treatment of Premature Ejaculation in the Asia-Pacific Region: Results From a Phase III Double-blind, Parallel-group Study of Dapoxetine. *Journal of Sexual Medicine*. 2009
- 102 Dresser MJ, Desai D, Gidwani S, Seftel AD, Modi NB. Dapoxetine, a novel treatment for premature ejaculation, does not have pharmacokinetic interactions with phosphodiesterase-5 inhibitors. *International Journal of Impotence Research*. 2006;**18**: 104-10.
- 103 Porst H, McMahon C, Althof S, Sharlip I, Bull S, Rivas DA. Baseline characteristics and treatment outcomes for men with acquired or lifelong premature ejaculation with mild or no erectile dysfunction: Integrated analysis of two phase III dapoxetine trials. *Journal of Sexual Medicine*. 2010.
- 104 Levine L. Evaluation of Withdrawal Effects with Dapoxetine in the Treatment of Premature Ejaculation (PE). *Poster presented at SMSNA 2006*.
- 105 Althof S, Levine S, Corty E, Risen C, Stern E, Kurit D. Clomipramine as a treatment for rapid ejaculation: A double-blind crossover trial of fifteen couples *Journal of Clinical Psychiatry*. 1995;**56**: 402-07.
- 106 Atmaca M, Kuloglu M, Tezcan E, Semercioz A. The efficacy of citalopram in the treatment of premature ejaculation: a placebo-controlled study. *International Journal of Impotence Research*. 2002;**14**: 502-5.
- 107 Goodman RE. An assessment of clomipramine (Anafranil) in the treatment of premature ejaculation. *Journal of International Medical Research*. 1980;**8**: 53-9.
- 108 Kara H, Aydin S, Yucel M, Agargun MY, Odabas O, Yilmaz Y. The efficacy of fluoxetine in the treatment of premature ejaculation: a double-blind placebo controlled study. *Journal of Urology*. 1996;**156**: 1631-2.
- 109 McMahon CG. Treatment of premature ejaculation with sertraline hydrochloride: a single-blind placebo controlled crossover study. *Journal of Urology*. 1998;**159**: 1935-8.
- 110 Waldinger MD, Hengeveld MW, Zwinderman AH. Paroxetine treatment of premature ejaculation: a double-blind, randomized, placebo-controlled study. *American Journal of Psychiatry*. 1994;**151**: 1377-9.
- 111 Waldinger MD. Premature ejaculation: definition and drug treatment. *Drugs*. 2007;**67**: 547-68.
- 112 Marangell L, Dennehy E, Wisniewski S, et al. Case-control analyses of the impact of pharmacotherapy on prospectively observed suicide attempts and completed suicides in bipolar disorder: findings from STEP-BD *Journal of Clinical Psychiatry*. 2008;**69**: 916-22.
- 113 Stone M, Laughren T, Jones ML, Levenson M, Holland PC, Hughes A, et al. Risk of suicidality in clinical trials of antidepressants in adults: analysis of proprietary data submitted to US Food and Drug Administration. *British Medical Journal*. 2009;**339**: 2880-90.
- 114 Mann J, Emslie G, Baldessarini R, et al. ACNP Task Force report on SSRIs and suicidal behavior in youth. *Neuropsychopharmacology*. 2006;**31**: 473-92.
- 115 Khan A, Khan S, Kilts R, Brown W. Suicide rates in clinical trials of SSRIs, other antidepressants, and placebo: analysis of FDA reports *American Journal of Psychiatry*. 2003;**160**: 790-92.

- 116 Khan A, Khan S, Kolts R, Brown WA. Suicide rates in clinical trials of SSRIs, other antidepressants, and placebo: analysis of FDA reports. *American Journal of Psychiatry*. 2003;**160**: 790-92.
- 117 Black K, Shea CA, Dursun S, Kutcher S. Selective serotonin reuptake inhibitor discontinuation syndrome: proposed diagnostic criteria. *Journal of Psychiatry and Neuroscience*. 2000;**25**: 255-61.
- 118 Kim SW, Paick JS. Short-term analysis of the effects of as needed use of sertraline at 5 PM for the treatment of premature ejaculation. *Urology*. 1999;**54**: 544-7.
- 119 McMahon CG, Touma K. Treatment of premature ejaculation with paroxetine hydrochloride as needed: 2 single-blind placebo controlled crossover studies. *Journal of Urology*. 1999;**161**: 1826-30.
- 120 Strassberg DS, de Gouveia Brazao CA, Rowland DL, Tan P, Slob AK. Clomipramine in the treatment of rapid (premature) ejaculation. *Journal of Sex and Marital Therapy*. 1999;**25**: 89-101.
- 121 Waldinger MD, Zwinderman AH, Olivier B. On-demand treatment of premature ejaculation with clomipramine and paroxetine: a randomized, double-blind fixed-dose study with stopwatch assessment. *European Urology*. 2004;**46**: 510-15.
- 122 Salonia A, Rocchini L, Sacca A, Pellucchi F, Ferrari M, Del Carro U, et al. Acceptance of and discontinuation rate from paroxetine treatment in patients with lifelong premature ejaculation. *Journal of Sexual Medicine*. 2009;**6**: 2868-77.
- 123 Berkovitch M, Keresteci AG, Koren G. Efficacy of prilocaine-lidocaine cream in the treatment of premature ejaculation. *Journal of Urology*. 1995;**154**: 1360-61.
- 124 Busato W, Galindo CC. Topical anaesthetic use for treating premature ejaculation: a double-blind, randomized, placebo-controlled study. *British Journal of Urology, International*. 2004;**93**: 1018-21.
- 125 Aversa A, Pili M, Francomano D, Bruzziches R, Spera E, La Pera G, et al. Effects of vardenafil administration on intravaginal ejaculatory latency time in men with lifelong premature ejaculation. *International Journal of Impotence Research*. 2009;**21**: 221-27.
- 126 McMahon CG, Stuckey B, Andersen ML. Efficacy of Viagra:Sildenafil Citrate in Men With Premature Ejaculation. *Journal of Sexual Medicine*. 2005;**2**: 368-75.
- 127 Salonia A, Maga T, Colombo R, Scattoni V, Briganti A, Cestari A, et al. A prospective study comparing paroxetine alone versus paroxetine plus sildenafil in patients with premature ejaculation. *Journal of Urology*. 2002;**168**: 2486-69.
- 128 McMahon CG, McMahon CN, Leow LJ, Winestock CG. Efficacy of type-5 phosphodiesterase inhibitors in the drug treatment of premature ejaculation: a systematic review. *British Journal of Urology, International*. 2006;**98**: 259-72.
- 129 Cavallini G. Alpha-1 blockade pharmacotherapy in primitive psychogenic premature ejaculation resistant to psychotherapy. *European Urology*. 1995;**28**: 126-30.
- 130 Fein RL. Intracavernous medication for treatment of premature ejaculation. *Urology*. 1990;**35**: 301-3.
- 131 Safarinejad MR, Hosseini SY. Safety and efficacy of tramadol in the treatment of premature ejaculation: a double-blind, placebo-controlled, fixed-dose, randomized study. *Journal of Clinical Psychopharmacology*. 2006;**26**: 27-31.
- 132 Salem E, Wilson S, Bissada N, Delk J, Hellstrom W, Cleves M. Tramadol HCL has promise in on-demand use to treat premature ejaculation. *Journal of Sexual Medicine*. 2008;**5**: 188-93.
- 133 Kwak TI, Jin MH, Kim JJ, Moon DG. Long-term effects of glans penis augmentation using injectable hyaluronic acid gel for premature ejaculation. *International Journal of Impotence Research*. 2008;**20**: 425-28.
- 134 Shi WG, Wang XJ, Liang XQ, Liu ZQ, Huang MJ, Li SQ, et al. [Selective resection of the branches of the two dorsal penile nerves for primary premature ejaculation]. *Zhonghua Nan Ke Xue*. 2008;**14**: 436-8.
- 135 Althof S. Psychological treatment strategies for rapid ejaculation: Rationale, practical aspects and outcome. *World Journal of Urology*. 2005;**23**: 89-92.
- 136 Semans J. Premature ejaculation. *Southern Medical Journal*. 1956;**49**: 352-58.
- 137 Hawton K. Treatment of sexual dysfunctions by sex therapy and other approaches. *British Journal of Psychiatry*. 1995;**167**: 307-14.
- 138 Donahey K, Miller S. Applying a common factors perspective to sex therapy. *Journal of Sex Education and Therapy* 2000;**25**: 221-30.
- 139 Mohr D, Bentler L. Erectile dysfunction: A review of diagnostic and treatment procedures. *Clinical Psychology Review*. 1990;**10**: 123-50.
- 140 Tang W, Ma L, Zhao L, Liu Y, Chen Z. Clinical efficacy of viagra with behavior therapy against premature ejaculation [Chinese]. *Zhonghua Nan Ke Xue*. 2004;**10**: 366-67.

- 141 Li P, Zhy G, Xu P, Sun J, Wang P. interventional effect of behavioral psychotherapy on patients with
premature ejaculation [Chinese]. *Zhonghua Nan Ke Xue* 2006;**12**: 717-19.
- 142 Yuan P, Dai J, Yang Y, Guo J, Liang R. A comparative study on treatment for premature ejaculation:
citalopram used in combination with behavioral therapy versus either citalopram or behavioral therapy
alone [Chinese]. *Chinnesse Journal of Andrology*. 2008;**22**: 35-38.
- 143 Steggall M, Fowler C, Pryce A. Combination therapy for PE: Results of a small-scale study. *Sex and
Relationship Therapy*. 2008;**23**: 365-76.
- 144 Abdo CH, Afif-Abdo J, Otani F, Machado AC. Sexual satisfaction among patients with erectile dysfunction
treated with counseling, sildenafil, or both. *Journal of Sexual Medicine*. 2008;**5**: 1720-26.
- 145 Aubin S, Heiman J, Berger R, Murallo A, Yung-Wen L. Comparing sildenafil alone vs. sildenafil plus brief
couple sex therapy on erectile dysfunction and couples' sexual and marital quality of life: A pilot study.
Journal of Sex and Marital Therapy. 2009;**35**: 122-43.
- 146 Melnik T, Abdo CH. Psychogenic erectile dysfunction: comparative study of three therapeutic
approaches. *Journal of Sex and Marital Therapy*. 2005;**31**: 243-55.
- 147 Althof S. Sex therapy in the age of pharmacotherapy. *Annual Review of Sex Research*. 2006: 116-32.
- 148 Althof S. Treatment of Rapid Ejaculation: Psychotherapy, Pharmacotherapy, and Combined Therapy. In:
Leiblum S, ed. *Principles and Practice of Sex Therapy (4th Edition)*. New York: Guilford Press; 2007.
- 149 Perelman M. Sex coaching for physicians: Combination treatment for patient and partner. *International
Journal of Impotence Research*. 2003;**15**: S67-74.
- 150 Perelman M. A new combination treatment for premature ejaculation. A sex therapist's perspective.
Journal of Sexual Medicine. 2006;**3**: 1004-12.
- 151 McMahon C, Abdo C, Incrocci L, Perelman M, Rowland D, Stuckey B, et al. Disorders of orgasm and
ejaculation in men. In: Lue T, Basson R, Rosen R, eds. *Sexual Medicine: Sexual Dysfunctions in Men and
Women (2nd International Consultation on Sexual Dysfunctions)*. Paris: Health Publications; 2004:409-
68.
- 152 Chia S. Management of premature ejaculation -- a comparison of treatment outcome in patients with
and without erectile dysfunction. *International Journal of Andrology*. 2002;**25**: 301-05.
- 153 Li X, Zhang SX, Cheng HM, Zhang WD. [Clinical study of sildenafil in the treatment of premature
ejaculation complicated by erectile dysfunction]. *Zhonghua Nan Ke Xue*. 2003;**9**: 266-9.
- 154 Sommer F, Klotz T, Mathers MJ. Treatment of premature ejaculation: A comparative vardenafil and SSRI
crossover study. *Journal of Urology*. 2005;**173**: 202:abstract 741.
- 155 Kirkegaard C, Faber J. The role of thyroid hormones in depression. *European Journal of Endocrinology*.
1998;**138**: 1-9.
- 156 Musselman DL, Nemeroff CB. Depression and endocrine disorders: focus on the thyroid and adrenal
system. *British Journal of Psychiatry Supplement*. 1996: 123-28.
- 157 Boneff AN. Topical treatment of chronic prostatitis and premature ejaculation. *International Urology and
Nephrology*. 1972;**4**: 183-6.
- 158 Brown AJ. Ciprofloxacin as cure of premature ejaculation. *Journal of Sex and Marital Therapy*. 2000;**26**:
351-52.
- 159 El-Nashaar A, Shamloul R. Antibiotic treatment can delay ejaculation in patients with premature
ejaculation and chronic bacterial prostatitis. *Journal of Sexual Medicine*. 2007;**4**: 491-96.
- 160 Patrick DL, Rowland D, Rothman M. Interrelationships among measures of premature ejaculation: The
central role of perceived control. *Journal of Sexual Medicine*. 2007;**4**: 780-88.
- 161 Althof S, Brock G, Rosen R, Rowland D, Aquilina J, Rothman M, et al. Validity of the patient-reported
clinical global impression of change as a measure of treatment response in men with premature
ejaculation. *Journal of Sexual Medicine*. 2010;**epub**.
- 162 American Academy of Family Physicians. Definition of Primary Care.
www.aafp.org/online/en/home/policy/policies/p/primarycare.html accessed 11/11/09
- 163 Annon J. *Behavioral Treatment of Sexual Problems: Brief Therapy*. Hagerstown, MD: Harper & Row;
1976.
- 164 Sadowsky R. The role of the primary care clinician in the management of erectile dysfunction. *Reviews
in Urology*. 2002;**4**: S54-63.
- 165 Reichman M. Optimizing referrals and consultations. *Family Practice Management*. 2007: 38-42.
- 166 Murray M. Reducing waits and delays in the referral process. *Family Practice Management*. 2002: 39-42.

Figure 1: Algorithm for the Management of PE (with permission of D. Rowland)¹




Table 1- Summary of PE Guideline Recommendations

Topic	Recommendation	Level of Evidence
Definition of Lifelong PE	A male sexual dysfunction characterized by ejaculation which always or nearly always occurs prior to or within about one minute of vaginal penetration, and the inability to delay ejaculation on all or nearly all vaginal penetrations, and negative personal consequences, such as distress, bother, frustration and/or the avoidance of sexual intimacy	1a
Definition of Acquired PE	There are insufficient published objective data to propose a new evidence-based definition of acquired PE, although it is believed the proposed criterion for lifelong PE might be applied to acquired PE as well	5d
Prevalence of PE	Statistical analysis of population-based data indicate that 1%-3% of men ejaculate in under 1 minute.	3d
Average Ejaculatory Latency	In multinational studies the median IELT is 5.4 minutes and decreased significantly with age. Median IELT may differ between countries.	2a
Quality of Life	Negative effects on quality of life and interpersonal difficulty related to their PE have been consistently reported by men and their partners	1a-3a
Etiology	The etiology of premature ejaculation is not known. To date, no biological factor has been shown to be causative in the majority of men with PE.	
Assessment	The committee agreed that there was inadequate evidence to recommend screening or case-finding for PE, either in a general population or in any sub-population. However, it is recommended that men with ED be screened for PE	5d
	It is recommended that clinicians utilize the screening questions in Table 3 and that clinicians take a medical and psychosocial history	5d
	Since patient self-report is the determining factor in treatment seeking and satisfaction, it has been recommended that self-estimation by the patient and partner of ejaculatory latency be routinely assessed in clinical practice when PE is present	2b
	The PEP or IPE are currently the preferred questionnaire measures for assessing PE, particularly in the context of monitoring responsiveness to treatment	2b
	For lifelong PE, a physical examination is highly advisable but not mandatory and should be conducted in most if not all patients	5d
	For acquired PE a targeted physical examination is mandatory to assess for associated/causal diseases such as ED, thyroid dysfunction or prostatitis	5d
Treatment	There is robust evidence to support the efficacy and safety of on-demand dosing of dapoxetine for the treatment of lifelong and acquired PE. It has been approved in some countries	1a
	There is robust evidence to support the efficacy and safety of off-label daily dosing of the SSRIs paroxetine sertraline, citalopram, fluoxetine, and the serotonergic tricyclic, clomipramine, and off-label on-demand dosing of clomipramine, paroxetine, and sertraline for the treatment of lifelong and acquired PE	1a
	There is good evidence to support the efficacy and safety of off-label on-demand topical anaesthetics in the treatment of lifelong PE	1b
	There is contradictory evidence to support the efficacy and safety of off-label on-demand or daily dosing of PDE-5 inhibitors in the treatment of lifelong PE in men with normal erectile function. Treatment of lifelong PE with PDE-5 inhibitors in men	4d

	with normal erectile function is not recommended and further evidence-based research is encouraged to further understand conflicting data	
	Treatment of PE with of daily α 1-adrenoceptor antagonists (Tramadol) cannot be recommended	2d
	There is modest evidence supporting the efficacy of psychological/behavioral interventions in the treatment of PE	2b
	Combining pharmacological and psychological/behavioral treatments may be especially useful in men with acquired premature ejaculation where there is a clear psychosocial precipitant or lifelong cases where the individual or couple's responses to PE are likely to interfere in the medical treatment and ultimate success of therapy	2a
	There is reliable evidence to support the treatment of PE and co-morbid ED with ED pharmacotherapy. There is level 3c evidence to support the treatment of PE and co-morbid ED with ED pharmacotherapy in combination with PE pharmacotherapy	1a
	Selective dorsal nerve neurotomy or hyaluronic acid gel glans penis augmentation may be associated with permanent loss of sexual function and is not recommended in the management of PE	4
Outcome	Treatment outcome can be addressed in one simple, brief and validated question known as the Clinical Global Index of Change (CGIC). It asks patients, "Compared to before starting treatment, would you describe your premature ejaculation problem as: much worse, worse, slightly worse, no change, slightly better, better, or much better?"	1b

Table 2 – Definitions of Premature Ejaculation

Definition	Source
A male sexual dysfunction characterized by ejaculation which always or nearly always occurs prior to or within one minute of vaginal penetration, and the inability to delay ejaculation on all or nearly all vaginal penetrations, and negative personal consequences, such as distress bother, frustration and/or the avoidance of sexual intimacy	International Society of Sexual Medicine, 2008
Persistent or recurrent ejaculation with minimal sexual stimulation, before, on or shortly after penetration and before the person wishes it. The condition must also cause marked distress or interpersonal difficulty and cannot be due exclusively to the direct effects of a substance	DSM-IV-TR, 2000
For individuals who meet the general criteria for sexual dysfunction, the inability to control ejaculation sufficiently for both partners to enjoy sexual interaction, manifest as either the occurrence of ejaculation before or very soon after the beginning of intercourse (if a time limit is required, before or within 15 seconds) or the occurrence of ejaculation in the absence of sufficient erection to make intercourse possible. The problem is not the result of prolonged absence from sexual activity	International Statistical Classification of Disease, 10 th Edition, 1994
The inability to control ejaculation for a “sufficient” length of time before vaginal penetration. It does not involve any impairment of fertility, when intravaginal ejaculation occurs	European Association of Urology. Guidelines on Disorders of Ejaculation, 2001
Persistent or recurrent ejaculation with minimal stimulation before, on, or shortly after penetration, and before the person wishes it, over which the sufferer has little or no voluntary control, which causes the sufferer and/or his partner bother or distress	International Consultation on Urological Diseases, 2004
Ejaculation that occurs sooner than desired, either before or shortly after penetration, causing distress to either one or both partners	American Urological Association Guideline on the Pharmacologic Management of Premature Ejaculation, 2004
The man does not have voluntary, conscious control, or the ability to choose in most encounters when to ejaculate	Metz and McCarthy, 2003
The Foundation considers a man a premature ejaculator if he cannot control his ejaculatory process for a sufficient length of time during intravaginal containment to satisfy his partner in at least 50 percent of their coital connections	Masters and Johnson, 1970
Men with an IELT of less than 1 minute (belonging to the 0.5 percentile) have “definite” premature ejaculation, while men with IELTs between 1 and 1.5 minutes (between 0.5 and 2.5 percentile) have “probable” premature ejaculation (Figure 3). In addition, an additional grading of severity of premature ejaculation should be defined in terms of associated psychological problems. Thus, both definite and probable premature ejaculation need further psychological subclassification in nonsymptomatic, mild, moderate, and severe premature ejaculation	Waldinger et al, 2005b

Table 3 – Recommended and Optional Questions to Establish the Diagnosis of PE and Direct Treatment

Recommended Questions	What is the time between penetration and ejaculation (coming)?
For Diagnosis	Can you delay ejaculation? Do you feel bothered, annoyed and/or frustrated by your premature ejaculation?
Optional Questions-	When did you first experience premature ejaculation?
Differentiate Lifelong And Acquired PE	Have you experienced premature ejaculation since your first sexual experience on every/almost every attempt and with every partner?
Optional Questions -	Is your erection hard enough to penetrate?
Assess Erectile Function	Do you have difficulty in maintaining your erection until you ejaculate during intercourse? Do you ever rush intercourse to prevent loss of your erection?
Optional Questions	
Assess Relationship Impact	How upset is your partner with your premature ejaculation? Does your partner avoid sexual intercourse? Is your premature ejaculation affecting your overall relationship?
Previous Treatment	Have you received any treatment for you premature ejaculation previously?
Impact on Quality of Life	Do you avoid sexual intercourse because of embarrassment? Do you feel anxious, depressed or embarrassed because of your premature ejaculation?

Table 4 – Summary of Recommended Pharmacological Treatments for Premature Ejaculation

Drug	Daily Dose/ As Needed	Dose	IELT Fold Increase	Side Effects	Status	Level of Evidence
<i>Oral Therapies</i>						
Dapoxetine ^{19, 101}	As Needed	30-60mg	2.5-3	Nausea Diarrhea Headache Dizziness	Approved in some countries	1a
Paroxetine ¹¹⁰	Daily Dose	10-40mg	8	Fatigue Yawning Nausea Diarrhea Perspiration Decreased Sexual Desire Erectile Dysfunction	Off label	1a
Clomipramine ^{105, 107}	Daily Dose	12.5-50mg	6		Off label	1a
Sertraline ¹⁰⁹	Daily Dose	50-200mg	5		Off label	1a
Fluoxetine ¹⁰⁸	Daily Dose	20-40mg	5		Off label	1a
Citalopam ¹⁰⁶	Daily Dose	20-40mg	2		Off label	1a
Paroxetine ¹¹¹	Daily dose for 30 days and then As Needed	10-40mg	11.6		Off label	1a
Paroxetine ¹¹³	As Needed	10-40mg	1.4		Off label	1a
Clomipramine ¹²¹	As Needed	12.5-50mg	4	Off label	1a	
<i>Topical therapy</i>						
Lidocaine/Prilocaine ¹²⁴	As Needed	25 mg/gm Lidocaine 25 mg/gm Prilocaine	4-6.3	Penile numbness Partner genital numbness Skin irritation Erectile Dysfunction	Off label	1b

Appendix 1

Index of Premature Ejaculation (IPE)

These questions ask about the effects your sexual problems have had on your sex life over the past four weeks. Please answer the following questions as honestly and clearly as possible. In answering these questions, the following definitions apply:

- *Sexual intercourse* is defined as vaginal penetration (you entered your partner).
- *Ejaculation*: the ejection of semen from the penis.
- *Control*: ejaculating when you are ready
- *Distress*: meaning how frustrated, disappointed, or bothered you are by your premature ejaculation

Mark only one box per question.

1) ***Over the past four weeks***, when you had sexual intercourse, how often did you have control over when you ejaculated?

- No sexual intercourse (not applicable)
- Almost always or always
- More than half the time
- About half the time
- Less than half the time
- Almost never or never

2) ***Over the past four weeks***, when you had sexual intercourse, how much confidence did you have over when you ejaculated?

- No sexual intercourse (not applicable)
- High confidence
- Moderately high confidence
- Neither high nor low confidence
- Moderately low confidence
- Low confidence

3) *Over the past four weeks*, when you had sexual intercourse, how often was it satisfactory for you?

- No sexual intercourse (not applicable)
- Almost always or always
- More than half the time
- About half the time
- Less than half the time
- Almost never or never

4) *Over the past four weeks*, when you had sexual intercourse, how satisfied were you with your sense of control over when you ejaculated?

- No sexual intercourse (not applicable)
- Very satisfied
- Somewhat satisfied
- Neither satisfied nor dissatisfied
- Somewhat dissatisfied
- Very dissatisfied

5) *Over the past four weeks*, when you had sexual intercourse, how satisfied were you with the length of intercourse before ejaculation?

- No sexual intercourse (not applicable)
- Very satisfied
- Somewhat satisfied
- Neither satisfied nor dissatisfied
- Somewhat dissatisfied
- Very dissatisfied

6) *Over the past four weeks*, how satisfied have you been with your sex life overall?

- No sexual intercourse (not applicable)
- Very satisfied

- Somewhat satisfied
- Neither satisfied nor dissatisfied
- Somewhat dissatisfied
- Very dissatisfied

7) *Over the past four weeks*, how satisfied have you been with your sexual relationship with your partner?

- No sexual intercourse (not applicable)
- Very satisfied
- Somewhat satisfied
- Neither satisfied nor dissatisfied
- Somewhat dissatisfied
- Very dissatisfied

8) *Over the past four weeks*, how much pleasure has sexual intercourse given you?

- No sexual intercourse (not applicable)
- High pleasure
- Moderate high pleasure
- Neither high nor low pleasure
- Moderately low pleasure
- Low pleasure

9) *Over the past four weeks*, how distressed (frustrated) were you by how long you lasted before you ejaculated?

- No sexual intercourse (not applicable)
- Extremely distressed
- Very distressed
- Moderately distressed
- Slightly distressed

Not at all distressed

10) *Over the past four weeks*, how distressed (frustrated) have you been about your control over ejaculation?

No sexual intercourse (not applicable)

Extremely distressed

Very distressed

Moderately distressed

Slightly distressed

Not at all distressed

Premature Ejaculation Profile (PEP Items)

Over the past month, was your control over ejaculation during sexual intercourse:

Very poor Poor Fair Good Very good

Over the past month, was your satisfaction with sexual intercourse:

Very poor Poor Fair Good Very good

Over the past month, how distressed were you by how fast you ejaculated during sexual intercourse?

Not at all A little Moderately Quite a bit Extremely

Over the past month, to what extent did how fast you ejaculated during sexual intercourse cause difficulty in your relationship with your partner?

Not at all A little Moderately Quite a bit Extremely

Premature Ejaculation Diagnostic Tool (PEDT)

Patient Instructions

This next questionnaire to help identify men who may have a problem with ejaculating too soon during sexual activity. Even if you do not have difficulties, please answer all the questions:

Please mark X in the box that best represented your answer for each of the questions below.

Please mark only one box for each question.

While your experiences may change from time to time, please report your general experiences with intercourse.

Definition: Ejaculation here refers to ejaculation (release of semen) after penetration (when your penis enters your partner)

	Not difficult at all	Somewhat difficult	Moderately difficult	Very difficult	Extremely difficult
(1) How difficult is it for you to delay ejaculation?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
	Almost never or never 0%	Less than half the time 25%	About half the time 50%	More than half the time 75%	Almost always or always 100%
(2) Do you ejaculate before you wish?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
(3) Do you ejaculate with very little stimulation?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
(4) Do you feel frustrated because of ejaculating before you want to?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4
(5) How concerned are you that your time to ejaculation leaves your partner sexually unfulfilled?	<input type="checkbox"/> 0	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3	<input type="checkbox"/> 4